

# Solution Manual Exercise 1 Linear Algebra

## Exercise 2.2

It should be easy to see that  $A$  is given by

$$A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

Let us try to find what  $A$  looks like in the  $\{|+\rangle, |-\rangle\}$ -basis. We have

$$|0\rangle = \frac{|0\rangle + |1\rangle}{2} + \frac{|0\rangle - |1\rangle}{2} = \frac{|+\rangle + |-\rangle}{\sqrt{2}}$$

$$|1\rangle = \frac{|0\rangle + |1\rangle}{2} - \frac{|0\rangle - |1\rangle}{2} = \frac{|+\rangle - |-\rangle}{\sqrt{2}}$$

So  $A$  takes  $\frac{1}{\sqrt{2}}\begin{pmatrix} 1 \\ 1 \end{pmatrix}_{\pm}$  into  $\frac{1}{\sqrt{2}}\begin{pmatrix} 1 \\ -1 \end{pmatrix}_{\pm}$  and vice versa. It should be easy to see that

$$A = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}_{\pm}$$

does the job.

## Exercise 2.5

$$(1) \quad \left( |v\rangle, \sum_j \lambda_j |w_j\rangle \right) = \sum_i v_i^* \sum_j \lambda_j w_{ij} = \sum_j \lambda_j \sum_i v_i^* w_{ij} = \sum_j (|v\rangle, |w_j\rangle)$$

$$(2) \quad (|v\rangle, |w\rangle) = \sum_i v_i^* w_i = \sum_i w_i^{**} v_i^* = (|w\rangle, |v\rangle)^*$$

$$(3) \quad (|v\rangle, |v\rangle) = \sum_i v_i^* v_i = \sum_i |v_i|^2 \geq 0$$

We get equality if and only if all  $v_i = 0$  which means that  $|v\rangle = 0$

### Exercise 2.7

Two vectors are orthogonal if their inner product is zero.

$$\langle v | w \rangle = (1 \quad -1) \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 1 - 1 = 0$$

Normalized forms:

$$|w_N\rangle = \frac{|w\rangle}{\sqrt{\langle w | w \rangle}} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

$$|v_N\rangle = \frac{|v\rangle}{\sqrt{\langle v | v \rangle}} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$

### Exercise 2.9

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = |1\rangle\langle 0| + |0\rangle\langle 1|$$

$$\sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} = i|1\rangle\langle 0| - i|0\rangle\langle 1|$$

$$\sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} = |0\rangle\langle 0| - |1\rangle\langle 1|$$

### Exercise 2.11

The eigenvalues are denoted  $\lambda_1$  and  $\lambda_2$ , and their corresponding normalized eigenvectors are  $|v_1\rangle$  and  $|v_2\rangle$ .

$$\sigma_x : \quad \lambda_1 = 1, \quad \lambda_2 = -1, \quad |v_1\rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \quad |v_2\rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$

$$\sigma_y : \quad \lambda_1 = 1, \quad \lambda_2 = -1, \quad |v_1\rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ i \end{pmatrix}, \quad |v_2\rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -i \end{pmatrix}$$

$$\sigma_z : \quad \lambda_1 = 1, \quad \lambda_2 = -1, \quad |v_1\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad |v_2\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

The diagonal representation is given by  $\sigma = \sum_i \lambda_i |v_i\rangle \langle v_i|$

$$\begin{aligned}\sigma_x &= \frac{1}{2} \left[ (|0\rangle + |1\rangle) (\langle 0| + \langle 1|) - (|0\rangle - |1\rangle) (\langle 0| - \langle 1|) \right] \\ \sigma_y &= \frac{1}{2} \left[ (|0\rangle + i|1\rangle) (\langle 0| + i\langle 1|) - (|0\rangle - i|1\rangle) (\langle 0| - i\langle 1|) \right] \\ \sigma_z &= |0\rangle \langle 0| - |1\rangle \langle 1|.\end{aligned}$$

Note that all the sigma matrices can be written as

$$\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

when we use the corresponding eigenvectors as basis

### Exercise 2.17

Any normal matrix  $A$  has a spectral decomposition

$$A = \sum_i \lambda_i |i\rangle \langle i|$$

where  $\lambda_i$  are the eigenvalues and  $|i\rangle$  the corresponding eigenvectors. We then have

$$A^\dagger = \sum_i \lambda_i^* |i\rangle \langle i|$$

If then  $A$  is hermitian,  $A = A^\dagger$ , we have  $\lambda_i = \lambda_i^*$ . Then all  $\lambda$ s has to be real. The implication the other way is just as easy, when all eigenvalues of a normal matrix  $A$  are real we see from the spectral composition that  $A = A^\dagger$  so  $A$  is hermitian

### Exercise 2.18

Let  $|u\rangle$  be a eigenvector of  $U$  with eigenvalue  $\lambda_u$ . Then

$$U |u\rangle = \lambda_u |u\rangle \quad \text{and} \quad \langle u| U^\dagger = \langle u| \lambda_u^*$$

This gives

$$\langle u| U^\dagger U |u\rangle = \langle u| |\lambda_u|^2 |u\rangle = |\lambda_u|^2$$

But since  $U^\dagger = U^{-1}$

$$\langle u| U^\dagger U |u\rangle = 1$$

So  $|\lambda_u|^2 = 1$  and  $\lambda_u = e^{i\theta}$ .

### Exercise 2.24

Note that if a matrix  $T$  is Hermitian we have:

$$\langle v|T|v\rangle = \langle v|T^\dagger|v\rangle = (\langle v|T|v\rangle)^\dagger = (\langle v|T|v\rangle)^*$$

for any  $|v\rangle$ . This means that  $\langle v|T|v\rangle$  must be real for any Hermitian  $T$ . We then define

$$B = \frac{1}{2}(A + A^\dagger) \quad \text{and} \quad C = -\frac{i}{2}(A - A^\dagger).$$

It is easy to check that  $A = B + iC$ , and that  $B$  and  $C$  are Hermitian. If  $A$  is positive,  $\langle v|A|v\rangle \geq 0$  for any  $|v\rangle$ , we have

$$\langle v|A|v\rangle = \langle v|B|v\rangle + i\langle v|C|v\rangle = \beta + i\gamma$$

Since  $B$  og  $C$  are Hermitian,  $\beta$  and  $\gamma$  must be real. To keep  $\langle v|A|v\rangle$  positive we must have  $\gamma = 0$ . Since  $|v\rangle$  is arbitrary, this means that  $C$  must be zero, which gives  $A = A^\dagger$ , i.e.  $A$  is Hermitian.